

COMPUTATIONS OF FLOER HOMOLOGY FOR CERTAIN LAGRANGIAN TORI IN CLOSED 4-MANIFOLDS

ADAM C. KNAPP

ABSTRACT. We compute the Lagrangian Floer cohomology groups of certain tori in closed simply connected symplectic 4-manifolds arising from Fintushel - Stern knot/link surgery. These manifolds are usually not symplectically aspherical. As a result of the computation we observe examples where $HF(L_0) \cong HF(L_1)$ and L_0 and L_1 are smoothly isotopic but L_0, L_1 are not symplectically isotopic and are distinguished by $HF(L_0, L_1)$.

1. INTRODUCTION

For a pair L_0, L_1 of Lagrangian tori in a symplectic 4-manifold, there are several types of isotopy that we may consider:

- Smooth isotopy.
- Lagrangian isotopy. L_0 and L_1 are Lagrangian isotopic in X if there is an smooth isotopy of L_0 to L_1 through Lagrangians.
- Symplectic/Hamiltonian isotopy. These are Lagrangian isotopies of L_0 to L_1 which extend to isotopies of the ambient manifold X and preserve the symplectic structure. If we think of a smooth isotopy as given by integrating a time dependant vector field χ_t on X , then the condition for symplectic isotopy is that $\iota(\chi_t)\omega$ is closed and the condition for Hamiltonian isotopy is that $\iota(\chi_t)\omega$ is exact. Of course, in the case of a simply connected ambient manifold, all symplectic isotopies are Hamiltonian.

In [12], Vidussi shows that there are an infinite number of smoothly nonisotopic Lagrangian tori inside $E(2)_K$. In [3], Fintushel and Stern define an integer valued smooth invariant of these tori, distinguishing the smooth isotopy class of an infinite family. The examples of Lagrangian tori we consider here are of the same type found in [12] and [3]. That is, they occur in the “link portion” of the link surgery manifolds of Fintushel-Stern.

In [1] Eliashberg and Polterovich give examples of Lagrangian tori in $\mathbb{R}^4$ which are Lagrangian isotopic but not Hamiltonian isotopic. The examples considered here differ from these as they lie in closed manifolds and may be essential.

In [10], Seidel gives an infinite family of smoothly isotopic but symplectically nonisotopic Lagrangian spheres in an exact symplectic 4-manifold distinguished by their Lagrangian Floer homology. This result was extended in [11] to certain embeddings of these examples into $K3$ and Enriques surfaces. Seidel’s computation uses the “Morse-Bott” spectral sequence for clean intersections of Lagrangians from [9] to compute the Lagrangian Floer homology of the Lagrangian spheres. We

2000 *Mathematics Subject Classification.* Primary 53D12; Secondary 57R52.

Key words and phrases. Lagrangian tori, Lagrangian isotopy, Lagrangian Floer Homology.

Partially supported by NSF Grant DMS0305818 and DMS0244663 (FRG).

say L_0, L_1 have clean intersection if their intersection is a embedded submanifold and $T(L_0 \cap L_1) = TL_0 \cap TL_1$. The result of Pożniak in [9] was a chain homotopy between the Morse complex of the clean intersection and the Lagrangian Floer complex restricted to a neighborhood of the clean intersection. We proceed somewhat similarly here except that where in Seidel's examples the vanishing of higher order differentials comes for more or less for free from grading considerations, we show vanishing by an essentially topological argument.

I would like to thank Ronald Fintushel, Richard Hind, and Tom Parker for helpful conversations.

2. CONSTRUCTION

Vidussi's symplectic version of the Fintushel-Stern link surgery can be described as follows: Consider M_L , a 3 manifold obtained from zero surgery on a nontrivial fibered m component link L in S^3 with a fibration $\pi : M_L \rightarrow S^1$. Choose metrics on M_L and S^1 appropriately so that the fibration map π is harmonic. (Without loss of generality assume that the metric on S^1 gives it volume 1.) Then $S^1 \times M_L$ has a symplectic form $\omega = d\theta \wedge d\pi + *_3 d\pi$. Here θ is a coordinate on S^1 and $*_3 d\pi$ indicates the pullback of $*d\pi$ in M_L via the projection $S^1 \times M_L \rightarrow M_L$. The form ω is closed since π is harmonic and nondegenerate since L is fibered. If m_i are meridians to the components K_i of L , then $S^1 \times m$ is a symplectic torus of square zero. Let X_i be symplectic 4-manifolds each with a symplectic torus F_i of square zero and tubular neighborhood $N(F_i)$. Suppose that $\pi_1(X_i \setminus N(F_i)) = 0$, then the symplectic fiber sum

$$X_L = S^1 \times M_L \underset{F_i = S^1 \times m_i}{\#}_{i=1, \dots, m} X_i$$

is simply connected and symplectic. Symplectic and Lagrangian submanifolds of each X_i and of $S^1 \times M_L$ which do not intersect the $F_i = S^1 \times m_i$ remain so under this process. (We also note that on each link component the choice of meridian m_i does not matter since isotopies of m_i induce deformation equivalences of symplectic structures on X_L .)

Let γ be a loop on a fiber of π . Then with the specified symplectic form, $L_\gamma = S^1 \times \gamma$ is a Lagrangian torus in $S^1 \times M_L$. When γ and the m_i are disjoint, L_γ is also naturally a Lagrangian torus in X_L . It is this class of Lagrangian tori which we will be considering here.

There is one more observation of note. In the 3-manifold M_L there is a natural construction of a vector field μ , namely the vector field uniquely determined by $\iota(\mu)(*d\pi) \equiv 0$ and $\pi^*(d\text{vol}_{S^1})(\mu) \equiv 1$. By construction, the time t flow of μ preserves the fibers of π , moving them in the forward monodromy direction. Thus the time 1 flow of μ on M_L gives the monodromy map when restricted to a fiber of π . If we extend μ to a vector field on $S^1 \times M_L$ which we also call μ , then we note that $\iota(\mu)\omega$ is a closed, but not exact 1-form. Thus we get a 1-parameter family of symplectomorphisms ϕ_t on $S^1 \times M_L$ which are not Hamiltonian.

Consider the action of ϕ_t on our Lagrangian torus L_γ . When $t \notin \mathbb{Z}$, $\phi_t(L_\gamma) \cap L_\gamma = \emptyset$ as γ is moved to a disjoint fiber. However, when $t \in \mathbb{Z}$ it is possible that $\phi_t(L_\gamma)$ and L_γ intersect. Further, when the monodromy is of finite order, we can find a good choice of meridian m so that the symplectic isotopies of L_γ to its iterates under the monodromy stay away from the $S^1 \times m_i$. These symplectic isotopies survive as Lagrangian isotopies in X_L .

3. CALCULATION OF FLOER COHOMOLOGY

We use the variant of Lagrangian Floer cohomology over the universal Novikov ring Λ in [5]. In this theory, the construction of the Floer cohomology groups is defined when certain obstruction classes vanish. These classes count pseudoholomorphic discs with boundary on the Lagrangians. The following lemmas show that we are in the situation where these classes vanish and serve to compute the homology.

Lemma 3.1. *Suppose $(S^1 \times M_L, \omega)$ is as above and that the link L is nontrivially fibered in the sense that the genus of the fiber is at least 1. Then $S^1 \times M_L$ contains no pseudoholomorphic spheres. Further, suppose that γ_i , $i = 0, 1$, are loops on a fiber of π which meet transversely in exactly one point and let $L_i = L_{\gamma_i}$. Then all pseudoholomorphic discs in $S^1 \times M_L$ with boundary on L_0 or on L_1 are constant and there are no nonconstant Floer discs for L_0, L_1 .*

Note that we cannot extend this lemma to say that there are no pseudoholomorphic representatives of $\pi_2(S^1 \times M_L, L_0 \cup L_1)$. We see such a counterexample in Section 4.

Proof of Lemma 3.1. As was assumed, M_L is a fibration over S^1 with fiber Σ_g of genus $g \geq 1$ and projection π . As $[\gamma_0] \cdot [\gamma_1] = \pm 1$ in the homology of the fiber no nonzero multiples of the two may be homologous. Thus they represent distinct infinite order elements of $\pi_1(\Sigma_g, \gamma_0 \cap \gamma_1)$ for which no powers $i, j \neq 0$ give $[\gamma_0]^i = [\gamma_1]^j$. By considering the universal abelian cover $\Sigma_g \times \mathbb{R}$, we see that $\pi_1(\Sigma_g)$ injects into $\pi_1(M_L)$ by the inclusion of a fiber. Then the subgroups generated by γ_0, γ_1 intersect trivially in $\pi_1(M_L, \gamma_0 \cap \gamma_1)$.

Since the genus of the fiber $g \geq 1$, $\pi_2(M_L) = 0$ and thus $\pi_2(S^1 \times M_L) = 0$. Now consider the exact sequence

$$0 = \pi_2(S^1 \times M_L) \longrightarrow \pi_2(S^1 \times M_L, L_i) \longrightarrow \pi_1(S^1 \times \gamma_i) \xrightarrow{i} \pi_1(S^1 \times M_L)$$

It follows that $\pi_2(S^1 \times M_L, L_i) = \ker(i) = 0$ by our assumptions on γ_i . Therefore, the homomorphism $\int \omega$ on $\pi_2(S^1 \times M_L, L_i)$ given by choosing a representative and integrating the pullback of ω over the disc is the zero homomorphism.

Let $\Omega(L_0, L_1)$ denote the space of paths $\delta : ([0, 1], 0, 1) \rightarrow (S^1 \times M_L, L_0, L_1)$ and $\Omega_0(L_0, L_1)$ be that subset whose members are homotopic to a point. As $L_0 \cap L_1$ is connected, $\Omega_0(L_0, L_1)$ is also connected. Let i_0, i_1 be the inclusions of L_0 and L_1 into $S^1 \times M_L$ respectively. There is an evaluation map $p : \Omega_0(L_0, L_1) \rightarrow L_0 \times L_1$, $p(\delta) = (\delta(0), \delta(1))$. This is a Serre fibration whose fiber is homotopy equivalent to $\Omega_0(S^1 \times M_L, x)$. Thus $\pi_k(p^{-1}(\delta_0, \delta_1)) \cong \pi_{k+1}(S^1 \times M_L)$ and there is the exact sequence

$$\begin{array}{ccccccc} \pi_2(S^1 \times M_L) & \longrightarrow & \pi_1(\Omega_0(L_0, L_1)) & \xrightarrow{p_*} & \pi_1(L_0) \times \pi_1(L_1) & \xrightarrow{i_{0*} \cdot i_{1*}^{-1}} & \pi_1(S^1 \times M_L) \\ \downarrow \cong & & \downarrow = & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & \pi_1(\Omega_0(L_0, L_1)) & \longrightarrow & \mathbb{Z}^2 \times \mathbb{Z}^2 & \longrightarrow & \mathbb{Z} \times \pi_1(M_L) \end{array}$$

For the last map, the sequence is exact in the sense that

$$\text{im}(p_*) = \ker(i_{0*} \cdot i_{1*}^{-1}) = \{(a, b) \in \pi_1(L_0) \times \pi_1(L_1) \mid i_{0*}(a) \cdot (i_{1*}(b))^{-1} = e\}.$$

Then $\pi_1(\Omega_0(L_0, L_1)) \cong \ker(i_{0*} \cdot i_{1*}^{-1})$. Since the γ_i are nontrivial and nontorsion each i_{0*}, i_{1*} is individually injective on π_1 . So the kernel of $i_{0*} \cdot i_{1*}^{-1}$ depends only on the intersection of the images of i_{0*} and i_{1*} in $\mathbb{Z} \times \pi_1(M_L)$. Then as we know the subgroups generated by γ_0, γ_1 in $\pi_1(M_L)$ intersect trivially, we see that $\pi_1(\Omega_0(L_0, L_1)) \cong \mathbb{Z}$.

Consider $\mathbb{D}^2$ as the unit disc in $\mathbb{C}$ and let $\partial_+ = \partial\mathbb{D}^2 \cap \{z \in \mathbb{C} \mid \Re z \geq 0\}$ and $\partial_- = \partial\mathbb{D}^2 \cap \{z \in \mathbb{C} \mid \Re z \leq 0\}$. Consider $\mathbb{A}$ as the annulus $\{z \in \mathbb{C} \mid 1 \leq |z| \leq 2\}$ with boundary components $\partial_{|z|=1}$ and $\partial_{|z|=2}$. We may represent elements of $\pi_1(\Omega_0(L_0, L_1))$ as maps of the annulus $(\mathbb{A}, \partial_{|z|=2}, \partial_{|z|=1})$ into $(S^1 \times M_L, L_0, L_1)$. As with $\pi_2(S^1 \times M_L, L_i)$ there is a homomorphism on $\pi_1(\Omega_0(L_0, L_1))$ which we shall call $\int \omega$ which is given by integrating the pullback of ω over (in this case) the annulus. We now show $\int \omega$ on $\pi_1(\Omega_0(L_0, L_1))$ is the zero homomorphism.

To see this, consider a certain generator for $\pi_1(\Omega_0(L_0, L_1)) \cong \mathbb{Z}$ represented by a map $u : (\mathbb{A}, \partial_{|z|=2}, \partial_{|z|=1}) \rightarrow (S^1 \times M_L, L_0, L_1)$ for which $u(\mathbb{A}) \subset L_0 \cap L_1$ and $\partial_{|z|=1}, \partial_{|z|=2}$ both map to $\pm$ the generator of $\pi_1(L_0 \cap L_1) = \mathbb{Z}$. Clearly $\int_{\mathbb{A}} u^* \omega = 0$. Then $\int \omega \equiv 0$ on $\pi_1(\Omega_0(L_0, L_1))$.

If we have topological Floer disc, that is, a map

$$u : (\mathbb{D}^2, \partial_-, \partial_+) \rightarrow (S^1 \times M_L, L_0, L_1),$$

then the images of $\pm i$ lie in $L_0 \cap L_1$. As $L_0 \cap L_1 \cong S^1$ is connected, we can connect the images of $\pm i$ by an arc $\gamma : ([-\frac{\pi}{2}, \frac{\pi}{2}], \{-\frac{\pi}{2}\}, \{\frac{\pi}{2}\}) \rightarrow (L_0 \cap L_1, u(-i), u(i))$. There are, of course, two ways of doing this but that will be immaterial. Then define $\tilde{u} : (\mathbb{A}, \partial_{|z|=2}, \partial_{|z|=1}) \rightarrow (S^1 \times M_L, L_0, L_1)$ by

$$\tilde{u}(z) = \begin{cases} u \circ \phi & \text{if } \Re z < 0 \\ \gamma(\theta) & \text{if } \Re z \geq 0, z = re^{i\theta}, -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \end{cases}$$

Where ϕ is a diffeomorphism from the interior of $D = \mathbb{A} \cap \{z \in \mathbb{C} \mid \Re z \leq 0\}$ to the interior of $\mathbb{D}^2$ whose extension to the boundary takes $D \cap \partial_{|z|=2}$ to ∂_- , $D \cap \partial_{|z|=1}$ to ∂_+ , the part of D lying on the positive imaginary axis to i and the part of D lying on the negative imaginary axis to $-i$.

Then $\int_{\mathbb{A}} \tilde{u}^* \omega = \int_{\mathbb{D}^2} u^* \omega$. Now, $\tilde{u}$ defines an element of $\pi_1(\Omega_0(L_0, L_1))$ so $\int_{\mathbb{D}^2} u^* \omega = \int_{\mathbb{A}} \tilde{u}^* \omega = 0$. Thus as all nonconstant pseudoholomorphic curves have positive symplectic area, there are no nonconstant Floer discs. $\square$

Lemma 3.2. *Suppose that $(S^1 \times M_L, \omega)$ and the L_i satisfy the assumptions of Lemma 3.1. Let m_i be meridians to each component K_i of L each away from the γ_i and (X_i, ω_{X_i}) be a collection of symplectic 4-manifolds each containing an embedded symplectic torus F_i of square zero. Then for any almost complex structure on the fiber sum manifold X_L which on each side of the fiber sum is sufficiently close to one for which the F_i or $S^1 \times m_i$ is pseudoholomorphic, all (perturbed) pseudoholomorphic discs in X_L with boundary on L_0 or on L_1 and all Floer discs for L_0, L_1 are constant.*

Note that if J_t is a loop of almost complex structures, starting at a J as described, which is contained within a small neighborhood of J , then J_t -pseudoholomorphic strips (Floer discs) can be considered as solutions to the perturbed pseudoholomorphic curve equations.

Proof of Lemma 3.2. In [8], Ionel and Parker construct a 6 dimensional symplectic manifold Z with a map to $\mathbb{D}^2$ so that over $\lambda \in \mathbb{D}^2 \setminus \{0\}$ the fiber is the symplectic sum X_L and over 0 the fiber is the singular manifold $S^1 \times M_L \bigcup_{i=1, \dots, m} X_i$. Each fiber $X_{L, \lambda}$ is canonically symplectomorphic to $S^1 \times M_L \setminus N(S^1 \times m_i) \bigcup_{i=1}^m X_i \setminus N(F_i)$ away from the fiber sum region.

Further, the almost-complex structure J_Z on Z is chosen so that in the singular fiber $X_{L, 0}$, $S^1 \times m_i$ is a pseudoholomorphic torus and so that the restriction of J_Z to each fiber is an almost complex structure. (The singular fiber is pseudoholomorphic in the sense that each inclusion of X and $S^1 \times M_L$ is pseudoholomorphic.)

Suppose that γ is a smooth embedded path in $\mathbb{D}^2$ which passes through zero and that $\{\lambda_n\}_{n=0}^\infty \rightarrow 0$ with $\lambda_n \in \gamma \setminus \{0\}$. Since the L_i are disjoint from the fiber sum region, there are Lagrangian submanifolds $\tilde{L}_i$ in Z for which in every fiber X_{L, λ_n} above λ_n , $\tilde{L}_i \cap X_{L, \lambda_n} = L_{i, n}$ which is mapped to L_i under the canonical identification.

Suppose that we have a family of pseudoholomorphic discs Σ_n in X_{L, λ_n} that have boundary in $\tilde{L}_i$ for each n and all have the same homology class in $H_2(X_L, \tilde{L}_i; \mathbb{Z})$ when it is identified with $H_2(X_{L, \lambda_n}, L_{i, n}; \mathbb{Z})$ via the bundle over γ . (The projection is a bundle map away from $\lambda = 0$.) Then the elements of this family each represent a homology class $[\Sigma_n]$ in $H_2(Z, \tilde{L}_i; \mathbb{Z})$. As Z is itself symplectic with a symplectic form ω_Z which restricts to each regular fiber as the symplectic form $\omega_{X_{\lambda_n}}$ obtained by fiber sum parameterized by λ_n , the integral of $\omega_{X_{\lambda_n}}$ over Σ_n within X_{λ_n} is equal to the integral of ω_Z over Σ_n . As Σ_n all represent the same homology class, the integrals must all be equal. Then since energy for pseudoholomorphic curves is the integral of the symplectic form, we have a bound on energy. Then by Gromov compactness for pseudoholomorphic curves with Lagrangian boundary conditions, these discs converge to a pseudoholomorphic curve $\hat{u}$ with image in the singular fiber $X_{L, 0} \cong S^1 \times M_L \bigcup_{i=1, \dots, m} X_i$ and boundary on $L_{i, 0} \cong L_i$. The domain C of $\hat{u}$ is then a collection of spheres and discs.

We next state Lemma 3.4 of [7] within our context. Note that here, ν is a perturbation of the $\bar{\partial}_J$ operator. (We suppress the perturbation elsewhere.)

Lemma 1 (Ionel, Parker). *Suppose that C is a smooth connected curve and $f : C \rightarrow S^1 \times M_L$ is a (J, ν) -holomorphic map that intersects $S^1 \times m$ at a point $p = f(z_0) \in S^1 \times m$. Then either*

- (1) $f(C) \subset S^1 \times m_i$ for some i or
- (2) *there is an integer $d > 0$ and a nonzero $a_0 \in \mathbb{C}$ so that in local holomorphic coordinates centered at p*

$$f(z, \bar{z}) = (p^i + O(|z|), a_0 z^d + O(|z|^{d+1}))$$

where $O(|z|^k)$ denotes a function which vanishes to order k at $z = 0$.

We note that no irreducible component of $\hat{u}$ is mapped entirely into any $S^1 \times m_i$ as there are no nonconstant holomorphic maps of spheres into tori and all discs have boundary on L_i away from each $S^1 \times m_i$. Part (2) of the lemma shows us that on each component of the domain, $\hat{u}$ meets the F_i as algebraic curves do. Thus we know that $\hat{u}$ intersects each $S^1 \times m_i$ at a finite number of points and that the image of each spherical or disc component of the domain of $\hat{u}$ will lie entirely in one of the X_i or in $S^1 \times M_L$.

Then our map $\hat{u}$ splits into two pseudoholomorphic maps $\hat{u}_1 : (C_1, \partial C_1) \rightarrow (S^1 \times M_L, L_i)$ and $\hat{u}_2 : C_2 \rightarrow \coprod_i X_i$. Here $C = C_1 \cup C_2$ with C_1 a collection of spheres and discs and C_2 a (possibly empty) collection of spheres so that $C_1 \cap C_2 \subset \hat{u}^{-1}(\bigcup_{i=1}^m S^1 \times m_i)$ are nodes of C .

Thus we have obtained $\hat{u}_1$, a pseudoholomorphic curve in $S^1 \times M_L$ with boundary on L_i . By our Lemma 3.1 this map must be constant. Thus the image of $\hat{u}_1$ is in L_i disjoint from the fiber sum region. Then since the images of $\hat{u}_2$ and $\hat{u}_1$ are connected, $\hat{u}_2$ must have empty domain. Therefore, $\hat{u}$ is constant.

This implies that for λ sufficiently small, all of the discs Σ_n must have been contained in $S^1 \times M_L$ disjoint from the fiber sum region. Therefore by Lemma 3.1, they were constant in the nonsingular fiber sum.

The proof follows identically if we consider pseudoholomorphic Floer discs for L_0, L_1 . □

Theorem 3.3. *Let $(S^1 \times M_L, \omega)$ and L_0, L_1 satisfy the conditions of Lemma 3.1. Then*

$$HF_{S^1 \times M_L}(L_0, L_0) \cong HF_{S^1 \times M_L}(L_1, L_1) \cong H^*(T^2) \otimes \Lambda$$

and

$$HF_{S^1 \times M_L}(L_0, L_1) \cong H^*(S^1) \otimes \Lambda.$$

Corollary 3.4. *Under the conditions of Theorem 3.3, the Lagrangian tori $L_i = S^1 \times \gamma_i$ are not Hamiltonian isotopic in $(S^1 \times M_L, \omega)$.*

Proof of Theorem 3.3. We begin by computing $HF(L_0, L_0)$. Since we have found in Lemma 3.1 that $\pi_2(S^1 \times M_L, L_0) = 0$, we have that

$$HF(L_0, L_0) \cong H^*(T^2) \otimes \Lambda$$

as in Floer's original work [4]. Similarly, $HF(L_1, L_1) = H^*(T^2) \otimes \Lambda$.

Now we consider $HF(L_0, L_1)$. As L_0, L_1 intersect cleanly, Proposition 3.4.6 of [9] implies that in some neighborhood $N(L_0 \cap L_1)$ of $L_0 \cap L_1$ the Floer complex and Morse complex for some Morse function $f : L_0 \cap L_1 \rightarrow \mathbb{R}$ coincide. This allows us to consider a slight modification of the action spectral sequence of [5].

The universal Novikov ring Λ can be written as the ring of formal sums $\sum_i a_i T^{\lambda_i} e^{n_i}$ with

- (1) $\lambda_i \in \mathbb{R}$ and $n_i \in \mathbb{Z}$
- (2) for each $\lambda^* \in \mathbb{R}$, $\#\{i | \lambda_i \leq \lambda^*\} < \infty$

Here the T^λ parameter will be used to keep track of the action of a pseudoholomorphic disc. The formula $\deg T^\lambda e^n = 2n$ determines a grading on Λ and we denote by Λ^k the homogeneous degree k part.

An $\mathbb{R}^+$ filtration on Λ is given by

$$F^\lambda \Lambda = \left\{ \sum_i a_i T^{\lambda_i} e^{n_i} \mid \lambda_i \geq \lambda \right\}$$

We can then get a $\mathbb{Z}$ filtration by picking some $\lambda^* \in \mathbb{R}^+$ and setting $\mathcal{F}^q \Lambda = F^{q\lambda^*} \Lambda$. The homogeneous elements of level q are then $\text{gr}_q(\mathcal{F}\Lambda) = \mathcal{F}^q \Lambda / \mathcal{F}^{q+1} \Lambda$.

Then as in Theorem 6.13 of [5] we have a spectral sequence $E_r^{p,q}$ with

$$E_2^{p,q} = \bigoplus_k H^k(L_0 \cap L_1; \mathbb{Q}) \otimes \text{gr}_q(\Lambda^{(p-k)})$$

Thus $E_2 \cong H^*(S^1) \otimes \Lambda$. We now want to see that all higher order differentials vanish. Recall that in Lemma 3.1 we showed that $\omega|_{\pi_2(S^1 \times M_L, L_0 \cup L_1)} \equiv 0$. Since the Hamiltonian perturbation may be chosen to be very small so that the local curves are of area less than λ^* , Lemma 3.1 shows that we have found all of the discs to be counted. Therefore, there are no higher order differentials and

$$HF(L_0, L_1) = H^*(S^1) \otimes \Lambda$$

□

Theorem 3.5. *Suppose (X_L, ω) and L_i satisfy the conditions of Lemma 3.2 and that for each L_i , the Maslov class $\mu : \pi_2(X_L, L_i) \rightarrow \mathbb{Z}$ takes only even values. Then in the fiber sum manifold X_L ,*

$$HF_{X_L}(L_0, L_0) \cong HF_{X_L}(L_1, L_1) \cong H^*(T^2) \otimes \Lambda$$

and

$$HF_{X_L}(L_0, L_1) \cong H^*(S^1) \otimes \Lambda.$$

Corollary 3.6. *Under the conditions of Theorem 3.5, the Lagrangian tori $L_i = S^1 \times \gamma_i$ are not symplectically isotopic in X_L .*

In section 4, We give examples which are Lagrangian isotopic.

Proof of Theorem 3.5. This is proved using the same methods as in the previous theorem. Lemma 3.2 ensures that the obstructions to defining a small perturbation Lagrangian Floer cohomology on the $L_i \subset X_L$ vanish and we make the following local (in J) computations:

- Using the action spectral sequence for $HF(L_0, L_0)$ we see that $E_2 = H^*(T^2) \otimes \Lambda$. By Lemma 3.2, we see that all the higher order differentials vanish for a small perturbation. In this case, the calculation of $HF(L_0, L_0)$ goes through as in the case of Theorem 3.3 and we find that $HF(L_0, L_0) \cong H^*(T^2) \otimes \Lambda$. Similarly, we get $HF(L_1, L_1) \cong H^*(T^2) \otimes \Lambda$.
- Using the action spectral sequence for $HF(L_0, L_1)$ we see that $E_2 = H^*(S^1) \otimes \Lambda$. By Lemma 3.2, we see that all the higher order differentials vanish. Therefore, the calculation of $HF(L_0, L_1)$ goes through as in the case of Theorem 3.3 and we find that $HF(L_0, L_1) \cong H^*(S^1) \otimes \Lambda$.

Now we show that the computed groups are invariant. Invariance is guaranteed as long as in 1-parameter families, there are no discs with boundary on L_0, L_1 of index -1 which bubble off.

Our restriction on the Maslov class ensures that there are no index -1 discs which appear in the boundary of 1-parameter families. Thus there is a continuation isomorphism and we have a well defined invariant of Hamiltonian isotopy in X_L . □

4. EXAMPLE

If we let L be the union of the right-hand trefoil knot K_1 and one of its meridians K_2 , the loops $\gamma_1, \gamma_2, \gamma_3$ in Figure 1 are all *freely* smoothly isotopic in M_L and meet transversely pairwise.¹ We select the fibration $\pi : M_L \rightarrow S^1$ so that the Seifert surface containing γ_i shown in Figure 1 is the fiber. The smooth isotopies of γ_1

¹Though they are freely isotopic, the proof of Lemma 3.1 shows that they are not equal in π_1 i.e. fixing a basepoint.




FIGURE 1. γ_i , $i = 1, 2, 3$ intersecting pairwise transversely (within the fiber) in single points

and γ_2 to a common curve are shown in Figures 4 and 5. In each of these figures going from (1) to (2) involves sliding over the 0-surgery on the meridian, going from (2) to (3) is an isotopy. To relate the end results we have the additional move of “twisting up the corkscrew” which takes the curves each (3) to the other. Note that this smooth isotopy is different than the Lagrangian isotopy which we will mention later.

Since the γ_i are smoothly isotopic, the Lagrangian tori $L_i = S^1 \times \gamma_i$ are smoothly isotopic. As Lagrangians do not have canonical orientations we neglect the orientations of the loops here.

Addressing the comment made after Lemma 3.1, we note that we can choose the almost complex structure J so that the Seifert surface Σ (a T^2) is pseudoholomorphic. Then choosing any pair γ_i, γ_j ($i \neq j$), $\Sigma \setminus (\gamma_i \cup \gamma_j)$ is a disc. This is however, not a Floer disc as it does not satisfy the correct boundary conditions.

For the left and right handed trefoils, the monodromy is of order 6 and in the basis A, B shown in Figure 2 is given by the matrix

$$\begin{bmatrix} 1 & 1 \\ -1 & 0 \end{bmatrix}$$

On the (positive/negative) Hopf link the monodromy is a (positive/negative) Dehn twist about a curve parallel to the components. The connect sum of fibered links is fibered with monodromy which splits around the connect sum region.

From this computation of monodromy, we see that γ_2 is $\pm$ the 2nd and 5th image of γ_1 under the monodromy map and that γ_3 is $\pm$ the 1st and 4th image of γ_1 . Finally, γ_1 is sent to $-\gamma_1$ under the third iteration of the monodromy. Thus, as the monodromy gives a symplectic isotopy (c.f. Section 2) in $S^1 \times M_L$ the tori L_1, L_2, L_3 are all symplectically isotopic. In X_L they are all Lagrangian isotopic as the fiber sum is taken away from the isotopy. Despite the existence of these




FIGURE 2. Basis for monodromy on trefoil

symplectic isotopies in $S^1 \times M_L$, Theorem 3.3 shows that the $L_i = S^1 \times \gamma_i$ are not Hamiltonian isotopic there.

Now we consider how Theorem 3.5 applies. That all the hypotheses of the theorem are satisfied, except that on the Maslov class, is clear. With the following lemma we see that the remaining condition is satisfied. Then Theorem 3.5 shows that the L_i are not symplectically isotopic in X_L .

Lemma 4.1. *For the Lagrangian tori L_i , we may choose $X_i = E(1)$, and the particular identification of F_i and $S^1 \times m_i$ so that $\mu_{L_i} : \pi_2(X_L, L_i) \rightarrow \mathbb{Z}$ is even.*

Proof. As L is a two component link with odd linking number, X_L is a homotopy $E(2)$ and thus is spin. See [2]. In fact $E(2)_L$ is $E(2)_{\text{Trefoil}}$. We shall write $E(2)_L := X_L$. For such a 4-manifold, the first Chern class is an even multiple of the fiber.


Note that μ_{L_i} factors through $\pi_2(E(2)_L, L_i) \rightarrow H_2(E(2)_L, L_i)$. As $H_1(E(2)_L) = 0$, the Meyer-Vietoris sequence gives that the group $H_2(E(2)_L, L_i)$ is generated by elements of $H_2(E(2)_L) / \langle L_i \rangle$ and relative classes with boundary spanning $H_1(L_i)$. The Maslov index of a class β , $\mu_{L_i}(\beta)$, $\beta \in H_2(E(2)_L, L_i)$, will change by an even amount whenever an element of $H_2(E(2)_L) / \langle L_i \rangle$ is added as $c_1(E(2)_L)$ is divisible by 2. Thus if we can find a pair of relative discs whose boundaries generate $H_1(L_i)$ and whose Maslov indices are even, we have shown that μ is even.

For $i = 1, 2$, we will choose the identification of F_i and $S^1 \times m_i$ so that

- (1) $\text{pt} \times m_i$ is identified with a vanishing cycle on F_i and
- (2) $S^1 \times \text{pt}$ is identified with the sum of two vanishing cycles on F_i whose boundaries meet once, transversely, in F_i .

Because $\pi_1(E(1) \setminus F_i) = 1$, we may select an elliptic fibration on $E(1)$ with nodal fibers having vanishing cycles a and b where a, b generate $\pi_1(F_i)$. With the decomposition $F_i = a \times b$, identify a with $\text{pt} \times m$ and $a + b$ with $S^1 \times \text{pt}$. This gives us the desired identification of F_i and $S^1 \times m_i$.

Each of the γ_i bounds a four times punctured disc D_{0, γ_i} in M_L where three punctures are meridians of K_1 and one is a meridian to K_2 . See Figure 3. By our

FIGURE 3. An isotope of γ_1 bounding meridians to K_1 and K_2

choice of fiber sum gluing, each meridian of K_1 bounds a vanishing disc on its $E(1)$ side of the fiber sum. Similarly, the meridian to K_2 bounds a vanishing disc on its $E(1)$ side of the fiber sum. Take three copies of the vanishing disc D_1, D_2, D_3 from the $E(1)$ fiber summed to $S^1 \times m_1$ and one vanishing disc D_4 of from the $E(1)$ fiber summed to $S^1 \times m_2$ and form $D_{\gamma_i} = D_{0, \gamma_i} \cup D_1 \cup D_2 \cup D_3 \cup D_4$.

Each γ_i 's Lagrangian framing relative to that induced by trivializing over D_{γ_i} is -2 and is given by a pushoff in the Seifert surface. See [3]. The framing coming from this disc is then $-1 - 1 - 1 - 1 - (-2) = -2$ and gives us $\mu_{\gamma_i}(D_{\gamma_i}) = -2$.

By our choice of gluing, the $S^1 \times \text{pt} \subset L_i$ is bounded by a pair of vanishing discs. This pair of vanishing discs intersects at one point on F and so can be smoothed to a disc $D_{S^1 \times \text{pt}}$ with relative framing -2 . For this loop the framing defect from the torus is 0 (given by pushoff in the monodromy direction) and so $\mu_{L_i}(D_{S^1 \times \text{pt}}) = -2$.

Thus we have found a basis on which μ is even, hence μ is even. $\square$


This example generalizes to similar links with $K_1 = T_{2, 2n+1}$ where we find many Lagrangian but not symplectically isotopic tori.


REFERENCES

- [1] Y. Eliashberg and L. Polterovich, *The problem of Lagrangian knots in four-manifolds*, in Geometric topology (Athens, GA, 1993), Vol. 2 of *AMS/IP Stud. Adv. Math.*, 313–327, Amer. Math. Soc., Providence, RI (1997).
- [2] R. Fintushel and R. Stern, *Knots, links, and 4-manifolds*, Invent. Math. **134** (1998), no. 2, 363–400.
- [3] ———, *Invariants for Lagrangian tori*, Geom. Topol. **8** (2004) 947–968 (electronic).
- [4] A. Floer, *Morse theory for Lagrangian intersections*, J. Differential Geom. **28** (1988), no. 3, 513–547.
- [5] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian Intersection Floer Theory - Anomaly and Obstruction*, at <http://www.math.kyoto-u.ac.jp/~fukaya/fooo.dvi>.
- [6] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), no. 2, 307–347.

- [7] E. Ionel and T. Parker, *Relative Gromov-Witten invariants*, Ann. of Math. (2) **157** (2003), no. 1, 45–96.
- [8] ———, *The symplectic sum formula for Gromov-Witten invariants*, Ann. of Math. (2) **159** (2004), no. 3, 935–1025.
- [9] M. Poźniak, Floer homology, Novikov rings and clean intersections, Ph.D. thesis, University of Warwick (1994).
- [10] P. Seidel, *Lagrangian two-spheres can be symplectically knotted*, J. Differential Geom. **52** (1999), no. 1, 145–171.
- [11] ———, *Graded Lagrangian submanifolds*, Bull. Soc. Math. France **128** (2000), no. 1, 103–149.
- [12] S. Vidussi, *Lagrangian surfaces in a fixed homology class: existence of knotted Lagrangian tori*, J. Differential Geom. **74** (2006), no. 3, 507–522.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824
E-mail address: `knappa@math.msu.edu`

FIGURE 4. Isotopy of γ_1

FIGURE 5. Isotopy of γ_2